

Morphology as a Basis for Taxonomy of Large Spirochetes
Symbiotic in Wood-Eating Cockroaches and Termites: *Pillotina* gen.
nov., nom. rev.; *Pillotina calotermitidis* sp. nov., nom. rev.;
Diplocalyx gen. nov., nom. rev.; *Diplocalyx calotermitidis* sp. nov.,
nom. rev.; *Hollandina* gen. nov., nom. rev.; *Hollandina*
pterotermitidis sp. nov., nom. rev.; and *Clevelandina*
reticulitermitidis gen. nov., sp. nov.

DAVID BERMUDEZ,^{1†} DAVID CHASE,^{2‡} AND LYNN MARGULIS^{1*}

Department of Biology, Boston University, Boston, Massachusetts 02215,¹ and Division of Surgical Oncology,
Department of Surgery, School of Medicine, University of California Davis Medical Center, Davis, California 95617²

The purposes of this paper are (i) to present a framework for the morphometric analysis of large uncultivable spirochetes that are symbiotic in wood-eating cockroaches and termites; (ii) to revive, in accordance with the rules of the *International Code of Nomenclature of Bacteria*, the names of three genera (*Pillotina*, *Diplocalyx*, and *Hollandina*) and three species (*Pillotina calotermitidis*, *Diplocalyx calotermitidis*, and *Hollandina pterotermitidis*) for the same organisms to which the names were originally applied, because these names were not included on the 1980 Approved Lists of Bacterial Names; and (iii) to formally propose the name *Clevelandina reticulitermitidis* for a new genus and species of spirochetes from the termite *Reticulitermes tibialis*. None of these genera and species has been cultivated either axenically or in mixed culture; hence, all are based on type-descriptive material.

Large spirochetes living under the anaerobic or microaerophilic conditions in intestines of wood-eating cockroaches and termites have never been cultivated (31). A new genus and species, "*Pillotina calotermitidis*," was proposed in 1967 by Hollande and Gharagozlou (10) for a spirochete from the termite *Calotermes praecox*. Another new genus and species, "*Diplocalyx calotermitidis*," was proposed for a separate morphotype from the hindgut of the termite *Calotermes flavicollis* by Gharagozlou (8) in 1968. A third genus and species, "*Hollandina pterotermitidis*" from the hindgut of the termite *Pterotermes occidentis*, was described by To et al. in 1978 (31). The genus "*Clevelandina*" has been informally proposed previously (26). Reports on these types of spirochetes have been reviewed by Breznak (6).

The work of Hollande and Gharagozlou is of particular interest because hollow tubules, 25 nm in diameter, were reported to be in the protoplasmic cylinder. Immunocytochemical investigations demonstrate the presence of antitubulin immunoreactivity in these large spirochetes (25). Further investigations of small cultivable spirochetes have resulted in the isolation of tubulinlike proteins (1). However, not all large spirochete morphotypes appear to have cytoplasmic tubules. Characterization of spirochetes with or without tubules requires distinctions among morphotypes.

We have examined the following four species of subterranean and dry-wood-eating termites by electron microscopy: *Pterotermes occidentis*, *Reticulitermes hesperus*, *Reticulitermes tibialis*, and *Incisitermes minor*. Our observations provide a framework for the morphometric analysis of

uncultivable spirochetes that are symbiotic in wood-eating termites and cockroaches.

In this paper we revive the genera "*Pillotina*," "*Diplocalyx*," and "*Hollandina*" and the species "*Pillotina calotermitidis*," "*Diplocalyx calotermitidis*," and "*Hollandina pterotermitidis*," in accordance with the rules of the *International Code of Nomenclature of Bacteria* (24), for the same organisms to which the names were originally applied, because they were not included on the 1980 Approved Lists of Bacterial Names (30). We formally propose the name "*Clevelandina reticulitermitidis*" for the type species of the new spirochete genus "*Clevelandina*" from the termite *Reticulitermes tibialis*.

MATERIALS AND METHODS

Sources of termite spirochetes. Termites were collected as indicated in Table 1. They were maintained in a laboratory, and the guts were removed for light and electron microscopic observations (31, 33).

Light microscopy. For observation of living material, freshly dissected guts were placed upon a glass microscope slide, covered with 1 or 2 drops of Trager solution A or Trager solution U (34), and pierced with a dissection needle. A cover glass was then sealed over the preparation with petroleum jelly.

Electron microscopy. For preparation of thin sections, entire termite guts were fixed with 3% glutaraldehyde in 0.025 M phosphate buffer at pH 6.8, postfixed with 1% osmium tetroxide in the same buffer, dehydrated in acetone, and embedded in Spurr resin by using the methods of Bloodgood (3) and Bloodgood and Fitzharris (4) as modified by To et al. (31) (2 to 5% tannic acid was added to the fixative for some of the samples and the impregnation time was extended to 24 h). Thin sections were stained first with 2% uranyl acetate and then with 2% lead citrate.

* Corresponding author.

† Present address: Center for Great Lake Studies, University of Wisconsin-Milwaukee, Milwaukee, WI 53204.

‡ Deceased.

TABLE 1. Sources and identification of North American termites used in this study

| Isopteran insect | Locality | Collected by: | Identified by: |
|--|-------------------------------------|----------------|----------------|
| Subterranean termites (Reticulitermitidae) | | | |
| <i>Reticulitermes hesperus</i> | San Diego, Calif. | L. Margulis | D. Chase |
| <i>Reticulitermes tibialis</i> | Joshua Tree National Monument, Nev. | D. Chase | D. Chase |
| Dry-wood-eating termites (Kalotermitidae) | | | |
| <i>Pterotermes occidentis</i> | Tucson, Ariz. | L. Margulis | W. Nutting |
| <i>Incisitermes (Kalotermites) minor</i> | Los Angeles, Calif. | D. Chase | D. Chase |
| <i>Incisitermes (Kalotermites) snyderi</i> | Naples, Fla. | S. F. Bermudes | P. Luykx |

Morphometrics. A total of 15 morphological criteria were measured (Table 2 and Fig. 1). The averages reported below represent measurements of all material from the original publications and from 3 to 20 examples of each morphotypes from the termites which we examined. Diameters (criterion 1) in oblong cells were measured at the narrowest and widest points enclosed by the cell membrane. In crenulated cells, measurements were made at the tips of the crenulations. The number of flagella (criterion 2) is reported as the number inserted at each end, approximated by the median number observed. The number of flagella observed could range from zero (beyond the most distal insertion) to twice the number of flagella inserted at one end, where, toward the center, flagella overlapped with the flagella from the other end. Previous descriptions (8, 10, 26, 31) have reported the maximum number observed in cross sections. The sillon (10) or groove (criterion 3) is an invagination of the outer membrane toward the inner membrane (Fig. 1B). This structure provides a reference point for the description of other structures. Crenulations (criterion 4) are conspicuous folds in the outer membrane (Fig. 1B). Criteria 5 through 7 deal with the presence of coatings on the inner and outer membranes of the gram-negative cell walls (i.e., the glycocalyx or sheath); the use of a ratio results in partial compensation for slightly oblique sections. When such surface layers are absent, the ratio of their thickness to the thickness of the membrane itself is zero. Absolute values are also given below. Spirochetes and protoplasmic cylinder cross sections of oblong shape were measured at both their widest (Fig. 1C) and narrowest points. The method for determining the ratio of the protoplasmic cylinder diameter to the overall cell

diameter (criterion 8) is illustrated in Fig. 1C. By using a ratio, comparisons could be made among spirochetes of different diameters. Criterion 9, the angle subtended by the flagella, is a measurement of the distribution of the flagella in the periplasm (Fig. 1D). Flagellar bundles (criterion 10) were deemed to be present if the angle subtended by the flagella was less than 180° and in areas where the flagella were not present, the outer membrane was in close apposition to the inner membrane (Fig. 1E). In such cases, a concavity was formed by the outer membrane near the junction between the flagellar bundle and the protoplasmic cylinder. An angle was formed by three points. Two points were formed by the intersection of a straight line with the outermost region of the outer membrane (Fig. 1E, line A, points 1 and 2). A third point was formed by the intersection of a line perpendicular to line A (line B) with the outer membrane at a distance maximal from line A. The angle of concavity was measured as the angle formed by the intersection of a line between points 1 and 3 (line C) and a line between points 2 and 3 (line D). The angle was less than 180° when a bundle was present. Cytoplasmic tubules (criterion 14) were small, hollow, walled structures as observed by electron microscopy. The polar organelle (synonym, polar membrane) (criterion 15) was an electron-dense lamina located on the inside of the inner membrane toward the distal end of the cell (Fig. 1B).

RESULTS AND DISCUSSION

Morphological characteristics. The major criteria (criteria 1 through 10) for distinguishing spirochete morphotypes are listed in Table 2. Criteria 11 through 15 provide less impor-

TABLE 2. Morphological criteria for assigning spirochetes to genera

| No. | Criterion Description | Figure illustrating criterion | Overall range in termite and cockroach spirochetes |
|-----|--|-------------------------------------|--|
| 1 | Diameter | 1A or C | 0.4–1.5 µm |
| 2 | Number of flagella | 1B | 15–70 |
| 3 | Presence of sillon | 1B | +, – ^a |
| 4 | Presence of crenulations | 1B | +, – |
| 5 | Ratio of thickness of outer coat of outer membrane to outer membrane thickness | 1B | 0–5 |
| 6 | Ratio of thickness of inner coat of outer membrane to outer membrane thickness | 1B | 0–12 |
| 7 | Ratio of thickness of outer coat of inner membrane to inner membrane thickness | 1B | 0–8 |
| 8 | Ratio of diameter of protoplasmic cylinder to cell diameter | 1C | 0.5–0.9 |
| 9 | Angle of protoplasmic cylinder subtended by flagella | 1D | 50–350° |
| 10 | Presence of flagellar bundle | 1E | +, – |
| 11 | Length | 1A | 10–>100 µm ^b |
| 12 | Amplitude | 1A | 0.3–0.6 µm ^b |
| 13 | Wavelength | 1A | 1.5–3.0 µm ^b |
| 14 | Presence of cytoplasmic tubules | 1B | +, – ^c |
| 15 | Presence of polar organelle | 1B | +, – |

^a +, Present; –, absent.

^b Not correlated with species.

^c The diameter of the tubules ranges from 15 to 25 nm.




FIG. 1. Morphological criteria for distinguishing spirochetes. The numbers in circles refer to the criteria listed in Tables 2 and 3. (A) Criteria visible by light microscopy (drawing by S. Alexander). (B) Idealized transverse section of *Pillotina* membranes and cytoplasmic tubules (drawing by J. Kaczman). (C) Two transverse sections showing how the ratio between the cell diameter and the periplasmic cylinder diameter is measured in irregularly shaped spirochetes (drawing by C. Lyons). (D) Three transverse sections indicating how angles subtended by flagella are measured (drawing by C. Lyons). (E) Transverse section showing how the angle of concavity is determined (drawing by C. Lyons).

tant but still useful bases for distinction among the morphotypes which we observed.

Spirochetes from the hindguts of wood-eating cockroaches and termites may possess a sillon, crenulations, or elaborate coatings of their membranes. The flagella and flagellar insertions of the large spirochetes have the appearance of the flagella and flagellar insertions of typical gram-negative bacteria (Fig. 2). The number of flagella ranges up to approximately 70 but is consistent within morphotypes. Some spirochetes tend to have flagella equally distributed around the protoplasmic cylinders, whereas in other spirochetes the flagella are confined to a limited portion of the circumference. Spirochetes with a low value for the angle subtended by the flagella tend to have their outer membrane tightly apposed to the inner membrane in regions where the flagella are not present. In those spirochetes with a high value (up to 350°) the outer membrane is not apposed, and the flagella are distributed throughout the periplasmic space. The thickness of the outer and inner membranes of the spirochetes which we examined ranged between 6.0 and 7.0 nm (average, 6.6 nm). Coatings on the outer and inner

membranes vary greatly in thickness among morphotypes but are relatively constant within individual morphotypes.

Length, amplitude, and wavelength are apparently consistent features of spirochetes observed by light microscopy. However, all of the termites examined harbor more than a single morphotype. The presence of morphotypes overlapping in diameter prevents the direct correlation of morphotype diameters observed by transmission electron microscopy with morphotype diameters observed by light microscopy. Because overall length, wavelength, and amplitude are not observable in individual transmission electron micrographs, diameter is the only morphological feature that is observable by both forms of microscopy. Thus, the spirochete features observed by transmission electron microscopy cannot yet be reliably correlated with any of the other morphological features observed by light microscopy. Because of this unresolved problem we disregarded previous reports on the length, amplitude, and wavelength (9, 10, 26, 31) of these spirochetes.

The presence of cytoplasmic tubules is more significant



FIG. 2. Flagellar insertions (arrows) of *Pillotina* sp. from *Reticulitermes hesperus*. f, flagella. Bar = 0.5 μ m. Magnification, $\times 79,800$. Previously published by Margulis et al. (26). Published by permission of Springer-Verlag.

than their absence; even in spirochetes such as "*Pillotina*," in which tubules have been observed many times, never does every section of the organism have them (Fig. 3). Micrographs that are of high enough quality and show appropriate sections (anterior and apposed to the flagellar basal insertion) reveal the polar organelle (Fig. 4). Although not mentioned as such, the polar organelle is clearly visible in the large spirochetes described by Grimstone in 1963 (9) and in the *Cristispira* described by Ryter and Pillot (29).


Spirochete morphotypes. The hypertrophied intestines of healthy members of any given termite species contain the same spirochete morphotypes. The great consistency of spirochete morphotypes (Fig. 5A and 6) in all healthy individuals of a given species, regardless of cast and age, demonstrates that these morphotypes are identifiable and distinct. Although in this paper we describe only large morphologically complex spirochetes, all of the termites examined also contained an abundance of smaller spirochetes (diameter, $<0.4 \mu$ m) (Fig. 6).

The large spirochetes, which are characterized by their size, lack of hooked ends, and habitat (the anaerobic or microaerophilic digestive tracts of animal hosts), belong to the family *Spirochaetaceae*. Cockroach and termite gut spirochetes differ from previously described genera. They are enteric in these insect hindguts and are intolerant of atmospheric oxygen. Thus, their habitat distinguishes them from *Spirochaeta*, spp., which are free-living, and from *Cristispira* spp., which are aerotolerant inhabitants of the crystalline styles of mollusks. Unlike *Borrelia*, the termite and roach spirochetes are not pathogenic (26). *Treponema* spp. cells (Table 3) tend to be small, usually 0.1 to 0.4μ m in diameter, and have relatively few flagella (<10 flagella), whereas the cells of the spirochetes which we examined tend to have diameters in excess of 0.4μ m and 15 to 70 flagella. Furthermore, the cockroach and termite hindgut spirochetes exhibit the following morphological features which are virtually absent from the other spirochete genera: crenulations, sillons, and elaborate coatings on their outer membranes. Their morphological complexity, coupled with their large size and number of flagella, definitively distinguishes the cockroach and termite hindgut spirochetes from other spirochetes.

Applying the criteria described above to the large spirochetes studied by electron microscopy from the guts of wood-eating cockroaches and termites yields at least four major morphotypes (Table 4). Examples of these morphotypes are shown in Fig. 2 through 7. For these morphotypes we revive the three genera and species described below and formally propose the new genus *Clevelandina* and the new species *Clevelandina reticulitermitidis*.

Description of *Pillotina* (ex Hollande and Gharagozlou 1968)
gen. nov., nom. rev. *Pillotina* (Pil.lo.ti'na. M.L. gen. *Pillotina*, in honor of J. Pillot, a French microbiologist). Symbiotic in the guts of wood-eating cockroaches and termites; probably anaerobic or microaerophilic. Helical cells are 0.6 to 1.5μ m in diameter. Stellate profile with approximately 30 to 70 flagella distributed throughout the periplasmic space as viewed in transverse section. Outer membrane crenulated (i.e., pleated or folded) (Fig. 1B, 2, 3A, and 5A and B). Outer coat of outer membrane either present or absent. One of the grooves between the ridges is deeper and narrower, forming the sillon. The type species is *Pillotina calotermitidis* from the termite *Calotermes praecox*.

Description of *Pillotina calotermitidis* (ex Gharagozlou 1968)
sp. nov., nom. rev. *Pillotina calotermitidis* (cal.o.ter.mit. id.is. M.L. gen. *calotermitidis*, named after the termite host *Calotermes* [synonyms, *Kalotermes*, *Postelectrotermes*] *praecox*). Helical cells are 1.0 to 1.5μ m in diameter, with 20 parallel ridges extending in a loosen spiral around the long axis nearly the full length, seen as crenulations in cross section. Alternate pairs of lower and higher ridges in the outer membrane are symmetrically placed on either side of the sillon. There are approximately 70 periplasmic flagella which are 15 to 18 nm in diameter. Angle subtended by the flagella, 350° ; concavity absent; no flagellar bundle. The distribution of the flagella is delimited by a lamina between the inner and outer membranes. An outer coat of the outer membrane is absent. The thickness of the inner coat of the outer membrane ranges from 22 to 77 nm (measured from the bottom of the crenulations toward the center of the cell) (average, 49.5 nm). The ratio of the thickness of the inner coat of the outer membrane to the outer membrane thickness ranges from 3.3 to 11.9 (average, 6.7). An outer coat of the inner membrane is absent. The protoplasmic cylinder is



circular to oblong in cross section and 0.5 to 0.7 μm in diameter. The ratio of the protoplasmic cylinder diameter to the cell diameter is 0.47 to 0.50 (average, 0.49). Microtubules, ca. 25 nm in diameter, are occasionally found within the protoplasmic cylinder. The type specimen (10) is from *Calotermes praecox* from the Island of Madeira, Portugal.

Description of *Diplocalyx* (ex Gharagozlou 1968) gen. nov., nom. rev. *Diplocalyx* (Dip.lo.ca.'lyx. Gr. *diploos*, twofold; L. n. *calyx*, cup, cover; M.L. n. *Diplocalyx*, twofold cover). Symbiotic in termites; probably anaerobic or microaerophilic. Helical cells are 0.7 to 0.9 μm in diameter. Approximately 40 to 60 periplasmic flagella are confined to a tight bundle between the inner and outer membranes, and there is a sill which varies in position in different morphotypes (26). A thick outer coat of the inner membrane, forming a calyx, is present except at the flagellar bundle and sill. Crenulations are lacking in the outer membrane. Microtubules, ca. 21 nm in diameter (Fig. 3B), and polar organelles are occasionally seen. The type species is *Diplocalyx calotermidis* from the termite *Incisitermes flavicollis*.

Description of *Diplocalyx calotermidis* (ex Gharagozlou 1968) sp. nov., nom. rev. *Diplocalyx calotermidis* (cal.o.ter.mi.'ti.dis. M.L. gen. *calotermidis*, named for the host termite *Calotermes* [synonyms *Kalotermes*, *Incisitermes*] *flavicollis*). Helical cells, lobate in cross section, are 0.8 μm in diameter, with approximately 40 flagella (diameter, 20 nm) in a tight bundle. The angle subtended by the flagella ranges from 50 to 70°. The angle of concavity ranges from 150 to 160°. The thickness of the outer coat of the outer membrane varies from 3.8 to 13.5 nm (average, 12 nm). The ratio of the thickness of the outer coat of the outer membrane to the outer membrane thickness is between 0.6 and 2.0 (average, 1.7). The thickness of the inner coat of the outer membrane ranges from 6 to 14 nm (average, 10 nm). The ratio of the thickness of the inner coat of the outer membrane to the inner membrane thickness varies from 0.9 to 2.1 (average, 1.5). The thickness of the outer coat of the inner membrane (the calyx) ranges from 42 to 51 nm (average, 48 nm). The ratio of the thickness of the outer coat of the inner membrane to the inner membrane thickness is between 6.4 and 7.7 (average, 7.3). The protoplasmic cylinder is slightly oblong and 0.38 μm (narrowest point) to 0.43 μm (widest point) in diameter. The ratio of the protoplasmic cylinder diameter to the cell diameter is between 0.48 and 0.53 (average, 0.51). The outer coat of the inner membrane is bisected by the sill. The position of the sill is opposite the flagellar bundle. The type specimen (8) is from *Calotermes flavicollis* found in southern France.

Description of *Hollandina* (ex To et al. 1978) Gen. nov., sp. nov., nom. rev. *Hollandina* (Hol.lan.din.'a. M.L. gen. *Hollandina*, in honor of Andre Hollande, Jr., a French protistologist). Symbiotic in the hindguts of wood-eating cockroaches and termites; probably anaerobic or microaerophilic. Helical cells rounded to oblong in cross section and 0.4 to 1.0 μm in diameter. A thick coat on the outer surface of the outer membrane is usually present, and

FIG. 3. Cytoplasmic tubules (arrows). Distribution and orientation vary. f, flagella. (A) *Pillotina* sp. from *Reticulitermes hesperus*. Tubules are 18 nm in both cross section and longitudinal section. c, crenulations. Bar = 0.5 μm . Magnification, $\times 52,200$. (B) *Diplocalyx* sp. from *Incisitermes minor*. Two tubules are 21 nm in diameter. s, sill. cx, calyx. Bar = 0.5 μm . Magnification, $\times 83,100$. (C) *Hollandina* sp. from *Incisitermes minor*. The cytoplasmic tubule is 20 nm in diameter. Bar = 0.5 μm . Magnification, $\times 79,800$.




FIG. 4. Polar organelle (P) of a *Hollandina* sp. from *Incisitermes minor*. Bar = 0.5 μ m. Magnification, $\times 52,600$. Previously published by Margulis et al. (26). Published by permission of Springer-Verlag.

crenulations and lacking. Cells have from 15 to 60 flagella distributed in the periplasmic space. A sillon is absent or present (Fig. 6). Polar organelles (Fig. 4) and microtubules (Fig. 3C) have been observed in some types. The type species is *Hollandina pterotermitidis* from the termite *Pterotermes occidentis*.

Description of *Hollandina pterotermitidis* (ex To et al. 1978) sp. nov., nom. rev. *Hollandina pterotermitidis* (pter.o. ter.mi.ti.'dis. M.L. gen. *pterotermitidis*, named for the host termite *Pterotermes occidentis*). Helical cells oblong in cross

section and 0.40 to 0.42 μ m (narrowest point) to 0.53 to 0.65 μ m (widest point) in diameter. The outer membrane is in close proximity to the inner membrane over a wide area for most of the cell length. A narrow groove (the sillon) is present toward the center. Cells have 15 flagella that are 15 to 16 nm in diameter. The angle subtended by the flagella ranges from 110 to 130° (average, 122°). The angle of concavity ranges from 170 to 180° (average, 175°). The thickness of the outer coat of the outer membrane ranges from 17 to 20 nm (average, 19 nm). The ratio of the thickness

TABLE 3. Comparative morphology of anaerobic symbiotic spirochetes: treponemes, brachyspiras, and borrelias

| Taxon | Length (μ m) | Width (μ m) | Wave-length (μ m) | Amplitude (μ m) | No. of flagella | Shape of ends ^a | Presence of cytoplasmic tubules | Presence of polar organelle | Presence of sheathed flagella | Presence of flagellar bundles | Reference(s) |
|---|-------------------|------------------|------------------------|----------------------|-----------------|----------------------------|---------------------------------|-----------------------------|-------------------------------|-------------------------------|--------------|
| Treponemes and brachyspiras (no vectors) | | | | | | | | | | | |
| <i>Treponema calligyrum</i> | 8–12 | 0.19–0.26 | 1.15 | 0.30 | 2–3 | ta | + ^b | NDT ^c | + | + | 16 |
| <i>Treponema genitalis</i> | 4–11 | 0.18 | 1.1 | 0.15 | 2–4 | ta | + | NDT | + | + | 15 |
| <i>Treponema hyodysenteriae</i> | 8–10 | 0.23–0.45 | 3.5 ^d | 0.5 ^d | 8–9 | ta | – ^d | NDT | + ^d | + ^d | 22 |
| <i>Treponema innocens</i> | 8–14 | 0.31–0.40 | 3.5 ^d | 0.5 ^d | 7–10 | ta | – ^d | NDT | + ^d | + ^d | 22 |
| <i>Treponema microdentium</i> | 4–10 | 0.15 | 0.9 | 0.15 | 1–3 | bl | + | NDT | + | + | 13 |
| <i>Treponema minutum</i> | 9–18 | 0.18 | 1.36 | 0.20 | 2–3 | ta | + | NDT | + | + | 13 |
| <i>Treponema pallidum</i> | 6–20 | 0.15 | 1.1 | 0.2–0.3 | 3 | pt | + | NDT | + | + | 12, 16 |
| <i>Treponema refringens</i> | 4–12 | 0.20–0.25 | 1.8 | 0.2–0.3 | 4–6 | ta | + | NDT | + | + | 17 |
| <i>Brachyspira aalborgi</i> | 1.7–6.0 | 0.2 | 2 | ND ^e | 4 | ta | – | NDT | + | + | 20 |
| Rumen spirochetes^f | | | | | | | | | | | |
| Strain PB (<i>Treponema saccharophilum</i>) | 12–20 | 0.6–0.7 | 2.4 | 0.54 | ≥ 32 | bl | ND | NDT | NDT | + | 27, 28 |
| Strain CA | 10–15 | 0.4–0.5 | ND | ND | 16–20 | ND | ND | ND | NDT | ND | 27 |
| Strain DA | 12–15 | 0.6 | ND | ND | ≥ 32 | ND | ND | ND | NDT | ND | 27 |
| Borrelias (vector transferred) | | | | | | | | | | | |
| <i>Borrelia baltazardi</i> | 14–20 | 0.35 | 2.0 | 0.35 | 25 | pt | – | NDT | – | + | 21 |
| <i>Borrelia burgdorferi</i> | 12–17 | 0.30 | 2.8 | ND | 7–11 | pt | – | NDT | + | + | 18 |
| <i>Borrelia merionesi</i> | 12–17 | 0.40 | 1.7 | 0.30 | 15–20 | pt | – | NDT | – | + | 14 |
| <i>Borrelia persica</i> | 16–23 | 0.45 | 2.0 | 0.35 | 25–30 | pt | – | NDT | – | + | 21 |
| <i>Borrelia recurrentis</i> | 12–16 | 0.50 | 1.7 | 0.30 | 15–20 | pt | – | NDT | – | + | 14 |

^a ta, Tapered; bl, blunt; pt, pointed.

^b +, Present; –, absent.

^c NDT, Not detected.

^d Unpublished data of K. Hovind-Hougen.

^e ND, No data available.

^f These anaerobic, host-associated spirochetes may require a new taxonomic treatment (28).




FIG. 5. *Pillotina* sp. from *Reticulitermes hesperus*. (A) Transverse sections showing crenulations (c), sillon (s), and consistency of ultrastructure. Bar = 0.5 μ m. Magnification, $\times 52,600$. (B) Longitudinal section. Bar = 0.5 μ m. Magnification, $\times 53,000$.




FIG. 6. *Hollandina* sp. from *Reticulitermes hesperus* showing consistency of ultrastructure and other small spirochetes (arrows). Bar = 0.5 μ m. Magnification, $\times 55,600$.

of the outer coat of the outer membrane to the outer membrane thickness is between 2.6 and 3.0 (average, 2.8). The thickness of the inner coat of the outer membrane ranges from 2 to 6 nm (average, 4.3 nm). The ratio of the thickness of the inner coat of the outer membrane to the outer membrane thickness is between 0.3 and 0.9 (average, 0.7). The thickness of the outer coat of the inner membrane

ranges from 7 to 13 nm (average, 9.5 nm). The ratio of the thickness of the outer coat of the outer membrane to the inner membrane thickness varies from 0.8 to 2.0 (average, 1.4). The oblong protoplasmic cylinder is 0.33 to 0.38 μ m (narrowest point) to 0.41 to 0.46 μ m (widest point) in diameter. The ratio of the protoplasmic cylinder diameter to the cell diameter ranges from 0.78 to 0.93 (average, 0.86).

TABLE 4. Distinctions among genera of large symbiotic spirochetes based on morphological criteria

| Criterion ^a | Genera | | | | | | |
|---------------------------------|---------------------------------|-----------------|-------------------|--------------------|------------------------|------------------------------|---|
| | <i>Cristispira</i> ^b | <i>Pilotina</i> | <i>Hollandina</i> | <i>Diplocatylx</i> | <i>Clevelandina</i> | <i>Borrelia</i> ^c | Rumen spirochetes (<i>Treponema</i>) ^d |
| Primary | | | | | | | |
| Diam (μ m) | 0.5–3.0 | 0.6–1.5 | 0.4–1.0 | 0.7–0.9 | 0.4–0.8 | 0.5 | 0.4–0.7 |
| Approximate no. of flagella | ≥ 100 | 30–70 | 30–60 | 40–60 | 30–45 | 15–20 | 8–16 |
| Presence of sillon | – ^e | + | +, – | + | + | – | – |
| Presence of crenulations | – | + | – | – | – | – | – |
| OCOM/OM ^f | ND ^g | 0 | 2.4–8.0 | 0.6–2.0 | 0–2.8 | 1–3 | ND |
| ICOM/OM ^h | ND | 3.3–11.9 | 0–1.3 | 0.9–2.1 | 2.8–5.0 (chambered) | 0 | ND |
| OCIM/IM ⁱ | ND | 0 | 1.0–2.5 | 6.4–7.7 | 0–3.6 | 1–2 | ND |
| PC/diameter ^j | 0.90 | 0.56–0.67 | 0.63–0.90 | 0.47–0.81 | 0.60–0.81 | 0.90–0.95 | 0.65–0.75 |
| Angle subtended by flagella (°) | 90–160 | 190–350 | 105–330 | 50–100 | 140–330 | 30–50 | 70–80 |
| Presence of flagellar bundles | + | – | +, – | + | +, – | – | + |
| Auxiliary | | | | | | | |
| Length (μ m) | 30–180 | ND | ND | ND | ND | 12–17 | 10–20 |
| Amplitude (μ m) | 4–6 | ND | ND | ND | ND | 0.3 | ND |
| Wavelength (μ m) | 10–20 | ND | ND | ND | ND | 1.7 | ND |
| Presence of cytoplasmic tubules | NDT ^k | + | + | + | NDT | – | NDT |
| Presence of polar organelle | + | + | + | + | NDT | – | NDT |

^a See Fig. 1.

^b See references 5, 23, and 29.

^c *Borrelia recurrentis* (16). Most other borrelias are smaller than 0.5 μ m in diameter.

^d Data from references 27 and 28. These anaerobic, host-associated spirochetes may require a new taxonomic treatment (28).

^e +, Present; –, absent.

^f Ratio of thickness of outer coat of outer membrane to outer membrane thickness.

^g ND, No data available.

^h Ratio of thickness of inner coat of outer membrane to outer membrane thickness.

ⁱ Ratio of thickness of outer coat of inner membrane to inner membrane thickness.

^j Ratio of diameter of protoplasmic cylinder to cell diameter.

^k NDT, Not detected.




FIG. 7. *Clevelandina* ultrastructure. s, sillon. cx, calyx. (A) *Clevelandina reticulitermitidis* from *Reticulitermes tibialis*, cross section. Bar = 0.5 μ m. Magnification, $\times 83,500$. (B) *Clevelandina* sp. from *Reticulitermes flavipes*, cross section. Bar = 0.5 μ m. Magnification, $\times 69,000$. Courtesy of J. Breznak. Previously published by Breznak (6) and Margulis et al. (26). Published by permission of The Williams & Wilkins Co.

The type specimen is from the hindgut of the termite *Pterotermes occidentis* from the Sonoran Desert in Arizona.

Description of *Clevelandina* gen. nov. *Clevelandina* (Cleveland, in 'a. M.L. gen. *Clevelandina*, in honor of L. R. Cleveland [1892-1969], an American biologist). Symbiotic in the hindguts of termites; probably anaerobic or microaerophilic. Helical cells are oblong in cross section and 0.4 to 0.8 μ m in diameter. Cells have 30 to 45 periplasmic flagella. The angle subtended by the flagella tends to be greater than 180°, and the inner coat (calyx) is correspondingly reduced. The cells are distinguished by a thick inner coat (layer) of the outer membrane with a chambered appearance and an outer coat of the inner membrane which is present except where the circumference is covered by flagella. Sillon present; no crenulations (Fig. 7A and B). The type species is *Clevelandina reticulitermitidis* from the termite *Reticulitermes tibialis*.

Description of *Clevelandina reticulitermitidis* sp. nov. *Clevelandina reticulitermitidis* (re.ti.cu.li.ter.mi.'ti.dis. M.L. gen. *reticulitermitidis*, named for the host termite genus *Reticulitermes*, in which it is found). Helical cells are slightly oblong and 0.52 to 0.61 μ m (narrowest point) to 0.67 to 0.73 μ m (widest point) in diameter. Approximately 45 flagella 17 to 20 nm in diameter are distributed throughout the periplasm. The angle subtended by the flagella is 260 to 270°; concavity absent; no flagellar bundle. The thickness of the outer coat of the outer membrane ranges from 0.0 to 18.6 nm (average, 11.5 nm). The ratio of the thickness of the outer coat of the outer membrane to the outer membrane thickness varies between 0.0 and 2.8 (average, 1.7). The inner coat of the outer membrane, which ranges from 25 to 27 nm thick (average, 25.3 nm), forms a chambered appearance in cross section. Chambers vary from 20 to 22 nm wide (average, 21 nm). The ratio of the thickness of the inner coat of the outer membrane to the outer membrane thickness varies from 3.8 to 4.1 (average, 3.9). The thickness of the outer coat of the inner membrane (calyx) varies between 12 and 20 nm (average, 14.6 nm). The ratio of the thickness of the outer coat of the inner membrane to the inner membrane thickness is between 1.8 and 3.8 (average, 2.8). The oblong protoplasmic




FIG. 8. Idealized drawings of *Treponema* and spirochete genera that are symbiotic in wood-eating cockroaches and termites (drawing by Laszlo Meszoly).

TABLE 5. Summary of distribution of spirochetes that are symbiotic in wood-eating cockroaches and termites^a

| Host family | Host genus | Location of host | Spirochete genus | Reference(s) |
|--|---|-----------------------|----------------------------------|--------------|
| Order Blattaria | | | | |
| Cryptocercidae (Problattidae; wood-eating cockroaches) | <i>Cryptocercus punctulatus</i> | United States | <i>Hollandina</i> | 10 |
| Order Isoptera | | | | |
| Hodotermitidae (damp-wood termites) | <i>Porotermes adamsoni</i> | Australia | — ^b | |
| Kalotermitidae (dry-wood termites) | <i>Bifiditermes condonensis</i> | Australia | — | |
| | <i>Ceratokaloterms apoliator</i> | Australia | — | |
| | <i>Cryptotermes brevis</i> | United States | — | |
| | <i>Cryptotermes cavifrons</i> | United States | — | |
| | <i>Cryptotermes gearyi</i> | Australia | — | |
| | <i>Glyptotermes iridipennis</i> | Australia | — | |
| | (<i>Kalotermes iridipennis</i>) | | | |
| | <i>Glyptotermes neotuberculatus</i> | Australia | — | |
| | <i>Kalotermes approximatus</i> ^c | United States | — | |
| | <i>Kalotermes banksiae</i> | Australia | — | |
| | <i>Kalotermes flavicollis</i> ^c | France, Spain | <i>Diplocalyx</i> | 8 |
| | <i>Kalotermes jouteli</i> | United States | — | |
| | (<i>Neotermes jouteli</i>) | | | |
| | <i>Kalotermes minor</i> | United States | <i>Hollandina</i> | 26 |
| | (<i>Incisitermes minor</i>) | | | |
| | <i>Kalotermes schwarzi</i> | United States | <i>Hollandina</i> | 26, 31 |
| | (<i>Incisitermes schwarzi</i>) | | <i>Pillotina</i> | 26, 31 |
| | <i>Kalotermes snyderi</i> ^d | United States | <i>Hollandina</i> ^e | |
| | <i>Incisitermes milleri</i> | United States | — | |
| | <i>Marginitermes hubbardi</i> | United States | <i>Diplocalyx</i> | 32 |
| | (<i>Kalotermes hubbardi</i>) | | | |
| | <i>Neotermes insularis</i> | Australia | — | |
| | <i>Paraneotermes simplicicornis</i> | United States | — | |
| | <i>Postelectrotermes praecox</i> | Madeira, Portugal | <i>Pillotina</i> | 10 |
| | (<i>Calotermes praecox</i>) ^c | | | |
| | <i>Pterotermes occidentis</i> | Mexico, United States | <i>Hollandina</i> | 31 |
| Mastotermitidae | <i>Mastotermes darwiniensis</i> | Australia | <i>Hollandina</i> | 7 |
| Rhinotermitidae (subterranean termites) | <i>Coptotermes aginaciformis</i> | Australia | — | |
| | <i>Coptotermes formosanus</i> | Hawaii | <i>Hollandina</i> | 32 |
| | <i>Heterotermes aureus</i> | United States | — | |
| | <i>Reticulitermes flavipes</i> | United States | <i>Clevelandina</i> | 6 |
| | | | <i>Pillotina</i> | 6 |
| | <i>Reticulitermes hesperus</i> | United States | <i>Clevelandina</i> ^f | |
| | | | <i>Hollandina</i> | 26 |
| | | | <i>Pillotina</i> | 26 |
| | <i>Reticulitermes tibialis</i> | United States | <i>Clevelandina</i> | 26 |
| | | | <i>Hollandina</i> | 26 |

^a See reference 26 for sources of termites unless otherwise noted.^b —, Absence of ultrastructural information on which identification could be based.^c Originally published nomenclature used (*Calotermes* is equivalent to *Kalotermes*).^d See Table 1.^e Unpublished data of B. Dorritie.^f Data not shown.

cylinder ranges from 0.34 to 0.42 μm (narrowest point) to 0.42 to 0.53 μm (widest point) in diameter. The ratio of the protoplasmic cylinder diameter to the cell diameter varies from 0.66 to 0.73 (average, 0.68). The type specimen (Fig. 7A) of *Clevelandina reticulitermitidis* is from *Reticulitermes tibialis* from Joshua Tree National Monument, Nev. (this termite was incorrectly identified as *Reticulitermes hesperus* by Margulis et al. [26]).

Distinctions among the genera. *Pillotina* is the only spirochete genus from termite or cockroach guts with crenulations. *Diplocalyx* has a flagellar bundle, sillon, and calyx. *Clevelandina* has a distinct chambered inner coat of the outer membrane. *Hollandina* tends to have thick coatings on the outer surface of the outer membrane and to lack crenu-

lations, a calyx, and a chambered coat. Idealized drawings of the genera are shown in Fig. 8.

Distribution of large symbiotic spirochetes. Large spirochetes are found associated with the digestive systems of wood-eating cockroaches and termites. The spirochetes from *Reticulitermes flavipes* (6) are clearly *Pillotina* sp. and *Clevelandina* sp. according to the criteria set forth in this paper. Other spirochetes described previously are also identifiably *Hollandina* sp. (e.g., the large surface spirochetes attached to the protist *Mixotricha paradoxa* from the Australian termite *Mastotermes darwiniensis* [7] and the unidentified spirochetes pictured by Grimstone [9]; also see reference 11). The geographical and host distributions of these large spirochetes are summarized in Table 5.

Microtubulelike structures. Eucaryotic microtubules are nearly invariant at 24 nm in diameter, except in some protists, where they approach 36 nm. However, cytoplasmic tubules in spirochetes (e.g., Fig. 3) and other procaryotes vary over a factor of at least four (Fig. 3). The diameter of the treponeme tubules may be as small as 6 nm (17, 19), whereas the diameter of *Pillotina* tubules may be as large as 25 nm (10). These differences in distribution, size, and morphology imply the existence of different types of cytoplasmic tubules in procaryotes. Some may be components of viral particles (25). The possible evolutionary importance of these tubules has recently been discussed (2).

ACKNOWLEDGMENTS

We are grateful to R. Bloodgood, B. Dorritie, K. Hovind-Hougen, B. Paster, and L. P. To who shared their unpublished data with us and to I. D. Gharagozlu for her original plates (8, 10). We are especially grateful to K. Hovind-Hougen for aid with our work; our study owes a great debt to her superb morphometric analyses of the smaller spirochetes. We thank F. Craft and W. Ormerod for aid in electron microscopy and photography. We are grateful for responses from the following investigators to whom queries about spirochete structure were made: R. Bloodgood, J. Breznak, E. Canale-Parola, E. Cox, B. Davis, E. P. Greenberg, C. Harwood, S. Holt, K. Hovind-Hougen, R. C. Johnson, L. K. Jurk, B. Paster, H. S. Pankratz, R. M. Smibert, S. Tamm, and M. Yamin. We thank S. F. Bermudes, G. Bosco, and L. Olendzenski for collecting *Incisitermes* (*Kaloterme*s) and *Neotermes* from Florida. We are grateful to P. Luykx for termite identification. We thank S. Alexander (Fig. 1A), J. Kaczman (Fig. 1B), C. Lyons (Fig. 1C through E), and L. Meszoly (Fig. 8) for the illustrations, as well as J. Breznak (Fig. 7B). We appreciate the excellent services of G. Fleischaker in preparation of the tables. We thank Joanne Chase for providing micrograph negatives and notes after the death of her husband.

This work was supported by grant NGR 004-025 from the National Aeronautics and Space Administration, (to L.M.), by the Departamento de Microbiología, Universidad Autónoma de Barcelona, Bellaterra, Spain, by the Lounsbery Foundation, by a National Science Foundation Graduate Fellowship (to D.B.), by the Boston University Graduate School, and by the Veterans Administration Hospital, Sepulveda, Calif. (for D.C.).

LITERATURE CITED

- Bermudes, D., S. P. Fracek, Jr., R. A. Laursen, L. Margulis, R. Obar, and G. Tzertzinis. 1987. Purification of tubulin-like protein from *Spirochaeta bajacaliforniensis*. *Ann. N.Y. Acad. Sci.* **503**:515-527.
- Bermudes, D., L. Margulis, and G. Tzertzinis. 1987. The prokaryotic origin of undulipodia: application of the panda principle to the centriole enigma. *Ann. N.Y. Acad. Sci.* **503**:187-197.
- Bloodgood, R. A. 1975. Biochemical analysis of axostyle motility. *Cytobios* **14**:101-120.
- Bloodgood, R. A., and T. P. Fitzharris. 1976. Specific association of prokaryotes with symbiotic flagellate protozoa from the hindgut of *Reticulitermes* and the wood-eating cockroach *Cryptocercus*. *Cytobios* **17**:103-122.
- Breznak, J. A. 1984. Genus II. *Cristispira*, Gross 1910, 1944^{AL}, p. 46-49. In N. R. Krieg and J. G. Holt (ed.), *Bergey's manual of systematic bacteriology*, vol. 1. The Williams & Wilkins Co., Baltimore.
- Breznak, J. A. 1984. Hindgut spirochetes of termites and *Cryptocercus punctulatus*, p. 67-70. In N. R. Krieg and J. G. Holt (ed.), *Bergey's manual of systematic bacteriology*, vol. 1. The Williams & Wilkins Co., Baltimore.
- Cleveland, L. R., and A. V. Grimstone. 1964. The fine structure of the flagellate *Mixotricha paradoxa* and its associated microorganisms. *Proc. R. Soc. Lond. B Biol. Sci.* **157**:668-683.
- Gharagozlu, I. D. 1968. Aspect infrastructural de *Diplocaelyx calotermitidis* nov. gen. nov. sp., spirochaetale de l'intestin de *Calotermes flavicollis*. *C.R. Acad. Sci. Ser. D* **266**:494-496.
- Grimstone, A. V. 1963. A note on the fine structure of a spirochete. *Q. J. Microsc. Sci.* **104**:145-153.
- Hollande, A. C., and I. D. Gharagozlu. 1967. Morphologie infrastructurale de *Pillotina calotermitidis* nov. gen. nov. sp. spirochaetale de l'intestin de *Calotermes praecox*. *C. R. Acad. Sci. Ser. D* **265**:1309-1312.
- Holt, S. C. 1978. Anatomy and chemistry of spirochetes. *Microbiol. Rev.* **42**:114-160.
- Hovind-Hougen, K. 1972. Further observations on the ultrastructure of *Treponema pallidum* Nichols. *Acta Pathol. Microbiol. Scand. Sect. B* **80**:297-304.
- Hovind-Hougen, K. 1974. The ultrastructure of cultivable treponemes. II. *Treponema calligyrum*, *Treponema minutum* and *Treponema microdentium*. *Acta Pathol. Microbiol. Scand. Sect. B* **82**:495-507.
- Hovind-Hougen, K. 1974. Electron microscopy of *Borrelia merrisoni* and *Borrelia recurrentis*. *Acta Pathol. Microbiol. Scand. Sect. B* **82**:799-809.
- Hovind-Hougen, K. 1975. The ultrastructure of cultivable treponemes. III. *Treponema genitalis*. *Acta Pathol. Microbiol. Scand. Sect. B* **83**:91-99.
- Hovind-Hougen, K. 1976. *Treponema* and *Borrelia* morphology, p. 7-37. In R. C. Johnson (ed.), *The biology of parasitic spirochetes*. Academic Press, Inc., New York.
- Hovind-Hougen, K. 1976. Determination by means of electron microscopy of morphological criteria of value for the classification of some spirochetes, in particular treponemes. *Acta Pathol. Microbiol. Scand. Sect. B Suppl.* **225**:1-40.
- Hovind-Hougen, K. 1984. Ultrastructure of spirochetes isolated from *Ixodes ricinus* and *Ixodes dammini*. *Yale J. Biol. Med.* **57**:543-548.
- Hovind-Hougen, K., and A. Birch-Andersen. 1971. Electron microscopy of endoflagella and microtubules in *Treponema* Reiter. *Acta Pathol. Microbiol. Scand. Sect. B* **79**:37-50.
- Hovind-Hougen, K., A. Birch-Andersen, R. Henrik-Nielsen, M. Orholm, J. O. Pedersen, P. S. Teglbjaerg, and E. H. Thaysen. 1982. Intestinal spirochetosis: morphological characterization and cultivation of the spirochete *Brachyspira aalborgi* gen. nov., sp. nov. *J. Clin. Microbiol.* **16**:1127-1136.
- Karimi, Y., K. Hovind-Hougen, A. Birch-Andersen, and M. Asmar. 1979. *Borrelia persica* and *B. baltazardi* sp. nov.: experimental pathogenicity for some animals and comparison of the ultrastructure. *Ann. Inst. Pasteur (Paris)* **130 B**:157-168.
- Kinyon, J. M., and D. L. Harris. 1979. *Treponema innocens*, a new species of intestinal bacteria and emended description of the type strain of *Treponema hyodysenteriae*, Harris et al. *Int. J. Syst. Bacteriol.* **29**:102-109.
- Kuhn, D. A. 1981. The genus *Cristispira*, p. 555-563. In M. P. Starr, H. Stolp, H. G. Truper, A. Balows, and H. G. Schlegel (ed.), *The prokaryotes: a handbook on habitats, isolation, and identification of bacteria*. Springer-Verlag, Berlin.
- Lapage, S. P., P. H. A. Sneath, E. F. Lessel, V. B. D. Skerman, H. P. R. Seeliger, and W. A. Clark (ed.). 1975. International code of nomenclature of bacteria. 1975 Revision. American Society for Microbiology, Washington, D.C.
- Margulis, L., L. P. To, and D. Chase. 1978. Microtubules in prokaryotes. *Science* **200**:1118-1124.
- Margulis, L., L. P. To, and D. G. Chase. 1981. The genera *Pillotina*, *Hollandina* and *Diplocaelyx*, p. 548-554. In M. P. Starr, H. Stolp, H. G. Truper, A. Balows, and H. G. Schlegel (ed.), *The prokaryotes: a handbook on habitats, isolation, and identification of bacteria*. Springer-Verlag, Berlin.
- Paster, B. J., and E. Canale-Parola. 1982. Physiological diversity of rumen spirochetes. *Appl. Environ. Microbiol.* **43**:686-693.
- Paster, B. J., and E. Canale-Parola. 1985. *Treponema saccharophilum* sp. nov., a large pectinolytic spirochete from the bovine rumen. *Appl. Environ. Microbiol.* **50**:212-219.
- Ryter, A., and J. Pillot. 1965. Structure des spirochetes. II. Etude du genre *Cristispira* au microscope optique et au électronique. *Ann. Inst. Pasteur (Paris)* **109**:552-562.

30. Skerman, V. B. D., V. McGowan, and P. H. A. Sneath (ed.). 1980. Approved lists of bacterial names. *Int. J. Syst. Bacteriol.* **30**:225–420.
31. To, L., L. Margulis, and A. T. W. Cheung. 1978. Pillotinas and hollandinas: distribution and behavior of large spirochetes symbiotic in termites. *Microbios* **22**:103–133.
32. To, L. P., and L. Margulis. 1978. Ancient locomotion: prokaryotic motility systems. *Int. Rev. Cytol.* **54**:267–293.
33. To, L. P., L. Margulis, D. Chase, and W. L. Nutting. 1980. The symbiotic microbial community of the Sonoran Desert termite: *Pterotermes occidentis*. *BioSystems* **13**:109–137.
34. Trager W. 1934. The cultivation of a cellulose-digesting flagellate, *Trichomonas termopsidis*, and of certain other termite protozoa. *Biol. Bull. (Woods Hole)* **66**:182–190.